

- ✓
- b) expressing said receptor or portion thereof under conditions permitting specific binding of said compound to said receptor or portion thereof;
 - c) exposing said cell to a sample suspected of comprising said compound; and
 - d) detecting the presence of any compound specifically bound to said receptor or portion thereof,

thereby determining whether said compound specifically binds to said receptor or portion thereof.

29

40.

The method according to claim 39, further comprising the steps of preparing a cell extract from the cell transfected with said nucleic acid molecule, isolating a membrane fraction of said cell extract, and contacting said sample with said membrane fraction under conditions permitting binding of the compound to said fraction.

30

41.

The method according to claim ²⁸39, wherein said detecting is performed by monitoring a change in the signaling activity of said CCR5 chemokine receptor or portion thereof.

31

42.

The method according to claim ²⁸39, wherein said detecting is performed by monitoring the acidification rate of said host cell.

32

43.

The method according to claim ³⁰41, wherein said detecting is performed by monitoring the level of intracellular calcium in said host cell.

33

44.

The method according to claim ³⁰41, wherein said detecting is performed by monitoring the stimulation of an intracellular cascade.

34

45.

The method according to claim ³⁰41, wherein said detecting is performed by monitoring the level of inositol triphosphate.

35

46.

The method according to claim ²⁸39, wherein said compound is an agonist of CCR5.

36

47.

The method according to claim ²⁸39, wherein said compound is an antagonist of CCR5.

37

48.

The method according to claim ²⁸39, wherein said at least one ligand is selected from the group consisting of RANTES, MIP-1 α , and MIP-1 β .

38

49.

The method according to claim ²⁸39, wherein said cell is selected from the group consisting of CHO-K1, HEK293, BHK21, and COS-7.

39

50.

The method according to claim 39, wherein said cell is exposed to said sample suspected of comprising said compound, in the presence of a ligand for the CCR5 receptor.

00939226-084401

a2
Ant